

Studying Elite Suppressors to Understand Mechanisms of Immune Control of HIV

by

Christopher W. Pohlmeier

A dissertation submitted to Johns Hopkins University in conformity with the requirements
for the degree of Doctor of Philosophy

Baltimore, Maryland
April 2016

ABSTRACT

Individuals who control HIV infection despite not being treated with antiretroviral therapeutics, deemed Elite Controllers or Suppressors (ES), represent a model for a functional cure. MHC class I alleles HLA-B*57 and HLA-B*27 have been associated with the ES phenotype, suggesting a genetic component. Accordingly, the role of CD8+ T cells in the ability to control infection is an ongoing field of investigation. ES have qualitatively greater CTL responses to HIV infection in *in vitro* and *ex vivo* assays. HIV often escapes from CTL pressure by mutating epitopes that are presented by MHC molecules. These CTL escape mutations can prevent CD8-mediated killing if the infected host cannot recognize these modified viral epitopes. Interestingly, escape variants are often found in the plasma of ES but not in integrated HIV-proviral genomes. In this thesis, I will present evidence that ES can suppress viral replication in CD4+ T cells that are infected with HIV variants harboring CTL escape mutations.

While ES have qualitatively more effective CD8+ T cells *ex vivo*, this does not preclude other possible factors from being critical to the phenotype. The frequency of latently infected CD4+ T cells in the periphery of ESs is significantly lower than in chronically infected individuals; this phenotype is not due to an inability for the CD4+ T cells to be infected. The amount of intracellular RNA in infected cell types has not been well characterized. We hypothesized that ongoing viral replication due to a lack of antiretroviral therapy may lead to increased baseline cell associated HIV-1 mRNA levels, which may allow for greater CTL-mediated clearance of HIV, in turn keeping the size of the HIV-infected CD4 population low. Here we characterize baseline and induced expression of cell-associated HIV-1 mRNA levels in CD4+ T cells from ES.

PhD Dissertation Referees for Christopher W Pohlmeier

Robert F Siliciano, MD PhD, Professor SOM (faculty sponsor)

Joel N Blankson, MD PhD, Associate Professor SOM (faculty sponsor)

TABLE OF CONTENTS

Acknowledgements	v
Abstract.....	ii
Introduction	1
Chapter 1: CD8+ T cells from HLA-B*57 elite suppressors effectively suppress replication of HIV-1 escape mutants.	3
ABSTRACT	3
BACKGROUND	4
RESULTS	5
DISCUSSION	10
CONCLUSION	13
MATERIALS AND METHODS	13
FIGURES	17
Chapter 2: Elite Suppressors Have Low Levels of Cell-Associated HIV-1 mRNA that Increase upon T cell Activation	24
ABSTRACT	24
BACKGROUND	25
RESULTS	25
DISCUSSION/CONCLUSION	27
MATERIALS AND METHODS	28
FIGURES	30
Significance and Future Directions.....	31
References	33
Curriculum Vitae	40

TABLE OF FIGURES

Figure 1.....	17
FITNESS COST OF CANONICAL HLA-B*57 ESCAPE MUTATIONS	
Figure 2.....	18
SUPPRESSION OF REPLICATION OF NL4-3 ESCAPE VARIANT VIRUSES	
Figure 3.....	19
INDIVIDUAL SUPPRESSION OF NL4-3 ESCAPE VARIANT VIRUSES	
Figure 4.....	20
INTRACELLULAR CYTOKINE STAINING OF ES CD8 ⁺ T CELLS	
Figure 5.....	21
INTRACELLULAR CYTOKINE STAINING OF CP CD8 ⁺ T CELLS	
Figure S1.....	22
THEORETICAL NONLINEAR REGRESSION CURVES PLOTTED WITH EACH ESCAPE VARIANT	
Figure S2.....	23
COMPARISON OF STIMULATION STATUS FOR SUPPRESSIVE FUNCTION	
Figure 6.....	30
HIV-MRNA BEFORE AND AFTER PMA AND IONOMYCIN STIMULATION	

ACKNOWLEDGEMENTS

In the existence that I experience, the double pendulum only has one starting condition. I am where I am by living the life I have lived, which is the only life I could have known. It is therefore of no surprise that things are how they are. And every situation in every corner of this existence has been critical in my arriving at my current circumstance, yes. No single step in my life was sufficient for placing me here today, yet every one was necessary. Nevertheless, my perception of fate is colored by the human condition; specific instances appear to have greater degrees of significance than others. While my thanks is ultimately senseless, the sentiment remains and can be appreciated to contain value.

Let's do this chronologically.

I would like to acknowledge my parents, Lynda and Phil. They have given me many tools and instilled in me the values that made me who I am. I continue to learn from them today, and hope to continue for many years to come.

I would like to acknowledge my siblings, more so my brother, Jeff. More tools and values, past learning and future learning.

I would like to acknowledge several influential teachers throughout my primary education, whom I will not acknowledge individually.

I would like to acknowledge my friends in high school, whom I will not acknowledge individually.

I would like to acknowledge my friends in college, whom I will not acknowledge individually.

I would like to acknowledge my undergraduate research mentors, Takanari and Tasuku.

I would like to acknowledge the administrative staff of my graduate program, whom I will not acknowledge individually.

I would like to acknowledge my friends in graduate school, whom I will not acknowledge individually.

I would like to acknowledge my graduate program mentors, Bob and Joel.

I would like to acknowledge my lab mentors, whom I will not acknowledge individually.

I would like to acknowledge my life partner, Sarah.

I would like to acknowledge my thesis committee members, whom I will not acknowledge individually.

I would like to acknowledge my lab mates, coworkers, and seronegative and seropositive donors, whom I will not acknowledge individually (somewhat for HIPPA-sake).

INTRODUCTION

ES are HIV-infected individuals who control viremia without antiretroviral therapy^{1,2,3,4}. Due to the very low level of viremia, these individuals almost never, if ever, transmit virus to a new host. Additionally, ES rarely lose control and progress to AIDS. Understanding the mechanism by which these individuals are able to control HIV infection could lead to the discovery of a functional cure for HIV. Early studies suggested that these individuals were merely infected with a defective HIV variant^{5,6}. While infection with replication deficient viral strains may lower the burden to control HIV infection, many ES are infected with fully replication competent strains^{7,8,9,10}.

Nevertheless, it is entirely possible that control is merely happenstance. ES may be the individuals who reside in one tail of a normal distribution centered around some median viral load¹¹. While this idea is difficult to disprove, there does exist, to some degree, a hierarchy of HLA alleles with respect to the ability to control viremia^{12,13}; this suggests that immune pressure is involved in control of viremia. Also, ES disproportionately possess HLA-B*57 and HLA-B*27 alleles compared to the general population, both seronegative and seropositive^{14,15,16,17,18,19,20}. This association of HLA molecules with control of viremia strongly suggests a genetic component to the ES phenotype, specifically a CD8-mediated mechanism of control.

Possessing protective HLA alleles is not sufficient for immune-mediated HIV control. Upon stimulation with HIV peptides, HIV-specific CD8+ T cells isolated from the periphery proliferate and generate cytokines; peptide stimulation of CD8+ T cells incurs a more robust phenotype in ESs than MHC-matched individuals who do not control^{18,21} (Chronic Progressors, or CPs). Additionally, coculturing CD8+ T cells isolated from the blood of ES with HIV-infected autologous CD4+ T cells results in greater inhibition of viral production than observed in CP counterparts²².

HIV evades this CTL pressure *in vivo* by generating mutations in MHC-restricted epitopes. These CTL escape mutations are either not processed, loaded, or recognized properly, and the ternary complex cannot form. CD4+ T cells from ES generally harbor few CTL escape mutations, whereas CPs often harbor several²³. Interestingly, escape variants can be found in the plasma of ES, and this does not necessarily lead to loss of control²⁴. In chapter 1, we will characterize the ability for CD8+ T cells to respond to stimulation with either wild type or escape variant HIV Gag peptides and suppress viral infection in CD4+ T cells infected with HIV variants that either do or do not contain CTL escape mutations.

Without antiretroviral therapy, the low level of plasma viremia observed in ES may be a result of ongoing replication. Antiretroviral therapy is believed to halt viral replication^{25,26}. Without ongoing replication, antigen cannot be presented to CD8+ T cells, and CTL responses will dissipate^{27,28}. If ES experience ongoing replication, CD8+ T cells will be continually primed for infection, improving the cytotoxic capacity of CD8+ T cells in a way that may not be translatable to CPs. It is poorly understood what the baseline levels of cell-associated HIV mRNA are and if these levels are due to ongoing replication. In chapter 2, we characterize cell-associated HIV mRNA in CD4+ T cells of ES.

CHAPTER 1: CD8+ T CELLS FROM HLA-B*57 ELITE SUPPRESSORS EFFECTIVELY SUPPRESS REPLICATION OF HIV-1 ESCAPE MUTANTS.

Pohlmeyer CW, Buckheit III RW, Siliciano RF, Blankson JN. *Retrovirology*. 2013, 10:152. 10.1186/1742-4690-10-152

ABSTRACT

Background

Elite Controllers or Suppressors (ES) are HIV-1 positive individuals who maintain plasma viral loads below the limit of detection of standard clinical assays without antiretroviral therapy. Multiple lines of evidence suggest that the control of viral replication in these patients is due to a strong and specific cytotoxic T lymphocyte (CTL) response. The ability of CD8+ T cells to control HIV-1 replication is believed to be impaired by the development of escape mutations. Surprisingly, viruses amplified from the plasma of ES have been shown to contain multiple escape mutations, and it is not clear how immunologic control is maintained in the face of virologic escape.

Results

We investigated the effect of escape mutations within HLA*B-57-restricted Gag epitopes on the CD8+ T cell mediated suppression of HIV-1 replication. Using site directed mutagenesis, we constructed six NL4-3 based viruses with canonical escape mutations in one to three HLA*B-57-restricted Gag epitopes. Interestingly, similar levels of CTL-mediated suppression of replication in autologous primary CD4+ T cells were observed for all of the escape mutants. Intracellular cytokine staining was performed in order to determine the mechanisms involved in the suppression of the escape variants. While low baseline CD8+ T cells responses to wild type and escape variant peptides were seen, stimulation of PBMC with either wild type or escape variant peptides resulted in increased IFN- γ and perforin expression.

Conclusions

These data presented demonstrate that CD8⁺ T cells from ES are capable of suppressing replication of virus harboring escape mutations in HLA-B*57-restricted Gag epitopes. Additionally, our data suggest that ES CD8⁺ T cells are capable of generating effective de novo responses to escape mutants.

BACKGROUND

In primary HIV-1 infection, vigorous viral replication results in plasma virus levels as high as one million copies/mL of HIV-1 RNA. As cellular immune responses develop, plasma virus levels decrease, and a viral set point is established. A subset of HIV-1-infected individuals known as Elite Controllers or Suppressors (ES), maintain a viral set point below the limit of detection of standard clinical assays (<50 copies of HIV-1 RNA/mL of blood^{1,2,3,4}). Early studies suggested that some long-term non-progressors (LTNPs) and ES are infected with defective viruses^{5,6}. In contrast, more recent studies have determined that many ES are infected with viruses that are fully replication competent, suggesting that host factors rather than infection by defective virus are responsible for ES status^{7,8,9,10,29}.

Some HLA alleles affect disease progression, including HLA-B*27, HLA-B*51, HLA-B*57/58, and HLA-B*35^{12,30}. Multiple cohort studies have demonstrated that the HLA-B*57 allele is overrepresented in ES^{14,15,16,17,18,19,20}. This finding has been confirmed by multiple GWAS studies^{31,32,33,34,35,36}. However, the majority of HIV-1-infected HLA-B*57⁺ patients develop progressive disease, and are thus termed chronic progressors (CPs)¹⁴. The protection conferred by HLA-B*57 and HLA-B*27 is thought to be mediated by effective CD8⁺ T cell responses against conserved immunodominant epitopes that are presented by MHC class I proteins^{18,21,22,37,38,39,40}. Comparison studies of HLA-B*57⁺ ES and CPs have provided some details about the elite suppressor phenotype. Stimulation of bulk peripheral blood mononuclear cells (PBMCs) with Gag peptides

induces greater proliferation as well as more robust granzymes A/B and perforin expression in CD8⁺ T cells from HLA-B*57⁺ ES compared to HLA-B*57⁺ CPs¹⁸.

Some nonsynonymous mutations in epitopes enable the virus to escape from the cytotoxic T lymphocyte (CTL) responses. The role these escape mutations play in determining protection versus progression in HLA-B*57 positive patients is controversial. A correlation between the number of HLA-B*57 Gag epitopes and the level of viremia was observed in a cohort of HLA-B*5703 positive patients with Clade C HIV-1 infection⁴¹. In another study, the development of escape mutations was temporally associated with virologic breakthrough in a patient who had maintained undetectable viral loads for a year after infection³⁴. In contrast, studies of early infection in other HLA-B*57/5801 positive patients have not found a correlation between the accumulation of escape mutations and virologic breakthrough^{43,44}. Furthermore, one study found no difference in the frequency of escape mutations in HLA-B*57-restricted epitopes in proviral clones amplified from ES when compared to CPs⁴⁵; other studies found that while escape mutations were largely absent from provirus amplified from CD4⁺ T cells of ES, virus amplified from the plasma of the same subjects contained a high frequency of escape mutations^{24,46}. In this study, we sought to explain how ES maintain undetectable levels of plasma virus despite the presence of circulating HIV-1 isolates that contain numerous escape mutations. Specifically, we asked whether CD8⁺ T cells from these patients were capable of inhibiting the viral replication of engineered escape mutants. We demonstrate that ES can inhibit the replication of escape variant HIV-1 and suggest that these patients are capable of generating protective *de novo* responses against the escape mutant variants. This work has implications for the design of therapeutic T cell vaccines to prevent the progression of HIV-1 disease.

RESULTS

Effect of escape mutations in HLA-B*57-restricted Gag epitopes on viral fitness

The effect of several escape mutations on viral fitness has been explored *in vivo* and *in vitro*^{47,48,49,50,51,52}. We focused on escape mutations in the three HLA-B*5703 restricted Gag epitopes: IW9 (Gag 147–155), KF11 (Gag 162–172), and TW10 (Gag 240–249). We have previously demonstrated that HLA-B*5703 positive ES in our cohort do not target the fourth HLA-B*57-restricted Gag epitope QW9 (Gag 308–316)²⁴ and therefore we did not include this epitope in our analysis. To address the fitness cost of these escape mutations, we introduced a series of mutations into the reference isolate NL4-3 and generated GFP-expressing HIV-1 pseudoviruses carrying these mutations. Virus concentrations were measured in triplicates by RT-PCR. We then infected PHA-activated CD4⁺ T cells from four seronegative healthy donors using a 2 log range of virus inoculums. Infection curves for the seven viruses are shown in Figure 1A.

Infectivity relative to the reference clone NL4-3 was taken as a measure of fitness.

Infectivity curves plateaued at different points for different mutants. A best fitting curve was generated with GraphPad to calculate a nonlinear least squares regression model and was used to determine the theoretical maximal infection. Figure S1 shows the best fit curve for each of the assayed viruses. This theoretical maximal infection was used to determine the relative percentage of maximal infection relative to the reference clone NL4-3 (Figure 1B). The I147L and A146P mutations, which are found in the IW9 epitope of p24, each had a minimal effect on viral fitness with calculated fitness levels of 88 percent and 90 percent respectively compared to unmutated NL4-3. The A163S mutation (72 percent relative fitness) had a larger effect on fitness consistent with a prior study that showed that mutations in this epitope in Clade C virus had a major impact on replication capacity⁴¹. When the A163S mutation was combined with the A146P mutation, a further decrease in fitness was observed (50 percent relative fitness). The

T242N/G248A variant had an intermediate level of fitness (67 percent relative fitness). Interestingly, the A146P/A163S/T242N/G248A variant (49 percent relative fitness) showed no additional loss of fitness than what was observed with the A146P/A163S variant.

ES CD8⁺T cells suppress replication of viruses containing CTL escape mutations

We next determined whether CD8⁺ T cells from ES could inhibit the infection of viruses harboring escape mutations. We isolated CD4⁺ and CD8⁺ T cells from seven HLA-B*5703⁺ ES and five healthy seronegative donors. The ES did not have any other protective alleles. Unstimulated CD4⁺ T cells were infected with each variant individually and were co-cultured with unstimulated CD8⁺ T cells at various effector to target ratios for three or five days. Interestingly, unstimulated CD8⁺ T cells from ES were capable of suppressing each variant (Figure 2A), whereas healthy donors had no suppressive capabilities (Figure 2D). The percentage of suppression decreased at lower effector to target ratios. Compared to day three, day five showed increased levels of suppression at all effector to target ratios (Figure 2A).

A prior study demonstrated that CD8⁺ T cells that were stimulated with peptides were more effective at eliminating HIV-1 infected target cells¹⁸. We therefore stimulated PBMCs from the same group of ES and healthy donors with a combination of three wild type peptides (wild type IW9, wild type KF11, and wild type TW10) or a combination of three escape variant peptides (mutant IW9 (I147L), mutant KF11 (A163S), or mutant TW10 (T242N/G248A)) in the presence of 10 units/mL IL-2. After seven days of stimulation, CD8⁺ T cells from both stimulation groups were individually isolated and cultured with unstimulated CD4⁺ T cells that were infected with either wild type virus or one of the 6 escape mutants. Interestingly, there was no statistical difference in the suppressive capacity of CD8⁺ T cells that were stimulated with either wild type or escape

variant peptides (Figure 2B, C and Figure S2). Figure 3 shows the suppressive ability of unstimulated CD8⁺ T cells from each patient at day three and day five. Statistically significant differences in levels of suppression between wild type and escape variants were seen only for A163P at a one to two effector to target ratio (Figure 3A); this difference was not seen at any other effector to target ratios. Taken together, these results demonstrate that ES are capable of recognizing escape variant epitopes as effectively as their non-mutated counterparts.

Stimulation of CD8⁺T cells with wild type or escape variant peptides increases expression of interferon γ and perforin

To determine the mechanism of CTL suppression of escape mutants, we analyzed the expression of IFN- γ and perforin in freshly isolated CD8⁺ T cells and CD8⁺ T cells that were primed with either wild type or escape variant peptides. Freshly isolated PBMCs and PBMCs from each stimulation group were stimulated overnight with individual wild type or escape variant peptides. Freshly isolated CD8⁺ T cells were observed to have very low levels of IFN- γ expression in response to each peptide (Figure 4A, B, C left panel). In contrast, culture of PBMCs with wild type peptides over a 7 day period prior to overnight stimulation with wild type peptides resulted in a statistically significant increase ($P < 0.05$) in IFN- γ expression (Figure 4A, B, C center panel). A similar response was seen when the CD8⁺ T cells were stimulated with the analogous escape variant peptide. Interestingly, IFN- γ expression in response to restimulation with wild type or escape variant peptides individually were similar in CD8⁺ T cells that were cultured for 7 days in the presence of either wild type or escape variant peptides (Figure 4A, B, C right panel).

Because perforin expression is associated with CTL-mediated killing^{18,39,53}, we also examined perforin expression by stimulated CD8⁺ T cells. Low levels of perforin and IFN- γ double positive cells were observed when freshly isolated CD8⁺ T cells were

stimulated with wild type or escape variant peptides (Figure 4D, E, F left panel). In contrast, CD8⁺ T cells that had been cultured for 7 days in the presence of wild type or escape variant peptide cocktails were observed to have an increase in the percentage of IFN- γ and perforin double positive CD8⁺ T cells when stimulated overnight with either wild type or escape variant peptides. Interestingly, culturing PBMCs in the presence of wild type peptides or escape variant peptide cocktails resulted in similar levels of CD8⁺ T cells that co-expressed IFN- γ and perforin in response to overnight peptide stimulation. Additionally, overnight stimulation with KK10 (Gag 263–272), an immunodominant HLA-B*27 specific peptide, resulted in no increase in IFN- γ or perforin expression, confirming the specificity of the enhanced CD8⁺ T cell responses to the HLA-B*57-restricted peptides.

To determine whether similar responses were present in HLA-B*57 positive CPs, we also analyzed CD8⁺ T cells from five CPs on suppressive HAART regimens, two CPs who recently had detectable viremia, and one CP with high levels of viremia who was not on HAART. With freshly isolated PBMCs, there was very low IFN- γ expression on CD8⁺ T cells in response to overnight stimulation with peptides (Figure 5A, B, C left panel). Stimulation of cells with wild type peptides or escape variant peptides induced an increase in IFN- γ expression by CD8⁺ T cells in some patients after overnight stimulation with TW10 or KF11 (Figure 5A, B, C center and right panels). The percentage of CD8⁺ T cells that co-expressed IFN- γ and perforin increased from the unprimed baseline (Figure 5E, F, G), though generally not as dramatically as the increase seen in ES. This is consistent with the enhanced proliferative capacity of ES HIV-specific CD8⁺ T cells^{18,21}. Interestingly, the patients on HAART who recently had detectable levels of viremia had higher responses than the patients on suppressive HART regimens who maintained undetectable viral loads consistent with a boosting effect of viral replication.

DISCUSSION

CTL responses against Gag epitopes have been associated with virologic control^{54,55,56} and Gag-specific CD8⁺ T cells can target incoming virions and therefore have the potential to kill cells prior to productive infection^{57,58,59,60,61}. The CTL response has been associated with the appearance of escape mutations in HIV and SIV infection. The mechanism of escape observed for individual mutations can vary. Mutations can affect epitope processing, stability of peptides, MHC:peptide complex stability, or TCR recognition of the MHC:peptide complex. Mudd *et al.* observed that Mamu-B*00801 macaques that controlled viral infection acquired few, if any, escape mutations in Vif and Nef epitopes, whereas macaques that progressed acquired several during the acute phase, suggesting that control may result from an immunologic pressure that prevents the appearance of escape mutations⁶². In contrast, Migueles *et al.* found that there was no difference between HLA-B*57⁺CPs and HLA-B*57⁺ ES in the frequency of escape mutations in Gag⁴⁵, and Bailey *et al.* found a high frequency of escape mutations in HLA-B*57-restricted epitopes present in virus amplified from ES plasma^{24,46}.

In this study, we sought to determine how ES maintain control of viral replication despite circulating escape mutant viruses in the plasma. We constructed a series of mutants that contained commonly observed HLA-B*57 restricted Gag escape mutations. While our study is limited by the fact that we did not study viral inhibition of autologous escape mutants isolated from each ES, the A146P and T242N/G248A mutations in the IW9 and TW10 epitopes are commonly seen in our cohort²⁴. Mutations in KF11 are rare in Clade B HIV-1 isolates, but one patient was found to have the A163S mutation and we demonstrated that this was in fact an escape mutation in a prior study²⁴. In agreement with other studies^{47,48,49,50,51}, we found that some of the escape mutants we generated were detrimental to viral fitness. While attenuating escape mutations may contribute to

elite control⁶³, viruses from CPs have been observed to have similar escape mutations, although compensatory mutations may partially restore viral fitness^{64,65}.

Klenerman and Zinkernagel demonstrated a limitation to the adaptive immune response: original antigenic sin⁶⁶. In brief, when CTLs respond to an intracellular pathogen, any variant of the original pathogen elicits the activation of the original memory response, which is potentially less effective in the face of the new variant of the pathogen⁶⁷.

Allen *et al.* demonstrated in a cohort of HLA-A*11⁺ individuals that the CTL response that recognizes escape variants was incapable of recognizing the original, un-mutated variants, as these CD8⁺ T cells express unique V β segments⁶⁸. New CTL responses have been shown to arise not only during the acute phase, but during chronic infection in HLA-A*02⁺ patients⁶⁹. Lichterfeld *et al.* have shown that HLA-B*27⁺ individuals can develop a *de novo* response to the immunodominant KK10 L268M escape mutation during chronic infection⁷⁰. In a previous study, HLA-B*57/58⁺ children infected perinatally showed a remarkable ability to generate *de novo* CD8⁺ T cell responses to escape mutations in the TW10 Gag epitope. Interestingly, there was little recognition of the wild type TW10 epitope in these children⁷¹. Another study found that CD8⁺ T cells from both HLA-B*57⁺ ES and HLA-B*57⁺ viremic patients made responses to autologous TW10 escape variant peptides⁷². Furthermore, we previously have described CD8⁺ T cell *de novo* responses to escape mutants in HLA-B*57⁺ES^{24,73}.

While of all these studies examined IFN- γ responses, secretion of this cytokine is not a correlate of immunity in HIV infection^{74,75}. Furthermore, discrepancies between IFN- γ ELISPOT assays and CD8⁺ T cell-mediated killing of both SIV and HIV escape variants have been reported^{76,77}. Therefore in order to determine whether protective *de novo* responses were present in ES, we looked at the ability of CD8⁺ T cells to suppress replication of escape mutants. We used the suppression assay because Saez-Cirion and

colleagues have demonstrated that the ability of unstimulated primary CD8⁺ T cells to inhibit viral replication correlates with elite control of HIV-1 infection^{78,80}, and we have recently confirmed this finding⁷⁹. In a prior study, we demonstrated that CD8⁺ T cells from an HLA-B*57 ES suppressed multiple rare autologous TW10 escape variants by a non-cross reactive *de novo* response⁸¹. In the current study, we demonstrated that this phenomenon is not limited to that one ES or to rare TW10 epitopes. CD8⁺ T cells from multiple HLA-B*5703⁺ ES were able to suppress the replication of virus containing common escape mutations in all three HLA-B*5703-restricted Gag epitopes. This is probably due to the development of CD8⁺ T cells that produce perforin in response to wild type and escape variant peptides.

Interestingly, while the presence of residual intracellular concentrations of antiretroviral drugs prevented us from infecting CD4⁺ T cells and performing the suppression assay with cells from HLA-B*57⁺ CPs, we demonstrated *de novo* IFN- γ production when CD8⁺ T cells of some CPs were stimulated with variant peptides. This is consistent with an earlier study which showed that HLA-B*57⁺ CPs made IFN- γ responses to autologous TW10 variants that were as strong, if not stronger, than the responses made by HLA-B*57⁺ ES⁷². Thus the elite control of viral replication is not solely due to the ability to recognize escape mutants. Rather, our work suggests that ES maintain control of viremia in spite of virologic escape in immunodominant epitopes because they develop protective CD8⁺ T cell responses to the escape variants. In contrast, CPs generally do not develop protective CTL responses^{18,21,39,53,78,79,80}, and a study that compared HLA-B*57⁺ ES to HLA-B*57⁺ CPs found that proliferative CTL responses as well as perforin secretion in response to HIV antigens correlated strongly with elite suppression²¹. Interestingly, we show here that stimulation with wild type and escape variant peptides can induce perforin responses to both peptides in some CPs. Thus, it

may be possible to immunize subjects with both wild type and escape variant peptides in order to induce protective CD8⁺ T cell responses that will prevent the emergence of common escape mutations. Taken together, it appears that ES CD8⁺ T cells may develop effective CTL suppressive responses to escape variants; these responses in addition to the reduced fitness of the escape variants, may explain how ES maintain levels of viremia in the face of virologic escape.

CONCLUSION

In this study, we demonstrate the ability of CD8⁺ T cells from ES to suppress replication of viruses harboring escape mutations in HLA-B*57-restricted Gag epitopes. The reduced fitness of these escape mutants may also contribute to elite control. Additionally, protective *de novo* CD8⁺ T cell responses to both wild type and escape variant peptides could be generated in ES and some CPs by priming PBMCs with either peptide. Induction of CD8⁺ T cells that could respond to wild type virus as well as common escape mutants would be advantageous for a CTL-based vaccine.

MATERIALS AND METHODS

Consent

All studies were approved by the Johns Hopkins Institutional Review Board. All patients and HIV negative donors provided written informed consent before participation in this study.

Construction of escape mutant viruses

Single-round infection by pseudotyped NL4-3 virus has been previously described⁸². In brief, *eGFP* was introduced in the *env* reading frame of pNL4-3, thus creating an *env* deficient pNL4-3 that allows for analysis of infected cells by flow cytometry. Individual point mutations were introduced by site directed mutagenesis (Agilent Technologies QuikChange II kit) and primers: A146P 5'

ggcaaatggtacatcagcccatatcacctagaactttaaatgc, I147L 5'

ggtacatcaggccctatcacctagaactttaaatgcatgg, A164S 5'

ggtaaaagtagtagaagagaaggtcttcagcccagaagtaatacc, and T242N/G248 5'

cccttcaggaacaaatagcgtggatgacacataatccacc followed by 5'

ggaactactagtaaccttcaggaacaaatagcgtgg. A146P/A163S was made by sequential mutagenesis. A146P/A163S/T242N/G248A was made by insertion of the digestion product of T242N/G248A with *SpeI* and *SbfI* into the A146P/A163S plasmid. These plasmids were sequence confirmed and individually cotransfected with pCI containing the III-B *env* reading frame into HEK293T cells using Lipofectamine 2000. 48 hours after transfection, supernatant was collected and virus was isolated by ultracentrifugation at $50,000 \times g$ through a 20% sucrose cushion for 2 h.

PBMC peptide stimulation

PBMCs were isolated from blood of ES, CP, and healthy donors by Ficoll gradient centrifugation. PBMCs were then cultured in RPMI 1640 supplemented with 10% FBS and 10 units/mL IL-2 in the presence of either 10 ug/mL total of TW10, KF11, and IW9 peptide (3 ug/mL each), or corresponding peptides containing escape mutations, for 7 days, with IL-2 supplemented every 48 h.

CD8⁺ And CD4⁺T cell isolation

PBMCs were isolated by Ficoll gradient centrifugation. CD8⁺ T cells were isolated from PBMC using the Miltenyi Human CD8 Microbeads according to the manufacturer's instructions. CD4⁺ T cells were isolated from either bulk PBMCs or CD8-depleted PBMCs using the Miltenyi Human CD4⁺ T Cell Isolation Kit II according to the manufacturer's instructions. Purity of both cell types was routinely greater than 95% as determined by staining with CD3-Pacific Blue and CD8 APC or CD4 PerCP-Cy5.5 (BD).

Suppression

Freshly isolated CD4⁺ T cells were spinoculated as described⁸³ at $1,200 \times g$ for 2 h with one of seven NL4-3 pseudotyped viruses in 2.9×10^6 cells/tube. We typically used 50 to 100 ng p24 virus for 100,000 CD4⁺ T cells which typically resulted in 2 to 10% GFP positive cells. Cells without virus were spinoculated as a negative control. After spinoculation, CD4⁺ T cells were washed and plated in a 96 well plate at 0.1×10^5 cells / well in RPMI 1640 supplemented with 10% FBS. Unstimulated, wild type stimulated, or Mutant stimulated CD8⁺ T cells were immediately added to the spinoculated CD4⁺ T cells, at specified ratios. Cells were cultured for 3 or 5 days before fixation and staining (CD3 Pacific Blue, CD8 APC, BD) and analysis by flow cytometry on a FACSCanto II (BD).

Intracellular cytokine analysis

0.5×10^6 unstimulated or stimulated PBMCs (from above) were restimulated with peptide (2 ug/mL), anti-CD28, and anti-CD49d in the presence of GolgiStop and GolgiPlug (BD) for 12 h. After staining with CD3 PE and CD8 APC-H7 (BD), cells were fixed and permeablized with Cytoperm/Cytofix Kit (BD). Cytokines were stained using IFN- γ PerCP-Cy5.5 (BD) and Perforin FITC (Cell Sciences). Stained cells were analyzed by flow cytometry on a FACSCanto II (BD).

Viral quantification

Concentrated virus was quantified by q-RTPCR. Viral RNA was isolated in triplicate with ZR Viral RNA Kit (Zymogen), and RT was performed using Superscript III First-Strand Synthesis Kit (Invitrogen) using poly(dT)₂₀ primers. VQA qPCR was performed as described⁸⁴. In brief, Taqman Fast Advanced Master Mix (Invitrogen) was used with primers 5' cagatgctgcatataagcagctg and 5' tttttttttttttttttttttgaagcac, and run on a ViiA7 (AB).

Fitness assay

PBMCs were isolated from healthy donors by Ficoll gradient centrifugation and activated in RPMI 1640 supplemented with IL-2 (100 units/mL) and PHA (1 ug/mL) for 3 days. CD4⁺ T cells were isolated (as above) and spinoculated ($1,200 \times g$, 2 h) in a 96-well plate with 0.1×10^6 cells / well, with different concentrations of each virus as shown in Figure 1. 72 h after spinoculation, cells were fixed (3.3% formaldehyde) and the level of infection for each concentration of virus was analyzed by flow cytometry on a FACSCanto II (BD). GraphPad Prism was used to generate model curves. Relative fitness was determined as a percentage of the maximal infection of an individual virus relative to the wild type control.

FIGURES




Figure 1

Fitness cost of canonical HLA-B*57 escape mutations. A. Average infection by seven NL4-3 escape variants in uninfected individuals. Viral inoculum was quantified by relative qPCR. 10^5 CD4⁺ T cells were infected in a 96-well plate in triplicate. Infection was determined by GFP expression by flow cytometry. Wild type NL4-3 (black) showed the highest level of infection, while NL4-3 escape mutant variants showed reduced maximal infection (I147L, orange; A146P, navy; A163S, red; A146P/A163S, purple; T242N/G248A, brown; A146P/A163S/T242N/G248A, teal). Error bars represent SEM. $n = 4$. **B.** Maximal infection of each escape variant is compared relative to wild type NL4-3 virus.




Figure 2

Suppression of replication of NL4-3 escape variant viruses. A-C: Unstimulated CD4⁺ T cells from HLA-B*5703 positive ES were infected with one of seven NL4-3 variants (wild type, black; I147L, orange; A146P, navy; A163S, red; A146P/A163S, purple; T242N/G248A, brown; A146P/A163S/T242N/G248A, teal) and cultured with autologous CD8⁺ T cells isolated from fresh PBMCs (**A**) or PBMCs stimulated with either wild type (**B**) or escape variant (**C**) HLA-B*57- Gag restricted Gag peptides for 7 days before isolation. CD8⁺ T cells were co-cultured with infected CD4⁺ T cells at three effector to target ratios. **D-F:** Unstimulated CD4⁺ T cells from healthy donors were infected with one of the seven NL4-3 variants used above and co-cultured with CD8⁺ T cells as was done with ES (**D**, unstimulated; **E**, wild type peptide stimulated; **F**, escape variant stimulated). Infection of CD4⁺ T cells was quantified by flow cytometry on days 3 (left) and 5 (right) after infection by GFP expression. Median values are plotted. For ES, n = 7. For healthy donors, n = 5.




Figure 3

Individual suppression of NL4-3 escape variant viruses. Unstimulated CD4⁺ T cells from HLA-B*5703⁺ ES were infected with one of seven NL4-3 variants (wild type, black; I147L, orange; A146P, navy; A163S, red; A146P/A163S, purple; T242N/G248A, brown; A146P/A163S/T242N/G248A, teal) and cultured with autologous unstimulated CD8⁺ cells at three effector to target ratios. **A** shows maximal suppression on day 3; **B** shows maximal suppression on day 5. Black horizontal bars indicate median. Asterisk indicates $P < 0.05$.




Figure 4

Intracellular cytokine staining of ES CD8⁺ T cells. **A-D:** CD8⁺ T cells of HLA-B*57 positive ES were either freshly isolated (left) or stimulated with either wild type (center) or escape variant (right) HLA-B*57-restricted Gag peptides for 7 days. Cells from each group underwent an overnight stimulation with individual peptides. Percentage of CD8⁺ T cells expressing IFN- γ when stimulated overnight with TW10 (**A**), KF11 (**B**), and IW9 (**C**) in blue, or the escape mutant variant peptide containing T242N/G248A (**A**), A163S(**B**), and I147L (**C**) mutations in red is shown. **E-H:** Percentage of CD8⁺ T cells expressing both IFN- γ and perforin after restimulation with TW10 (**E**), KF11 (**F**), and IW9 (**G**) in blue, or the escape mutant variant peptide containing T242N/G248A (**E**), A163S (**F**), and I147L(**G**) in red is shown. **D** and **H** show CD8⁺ T cells that express IFN- γ or co-express IFN- γ and perforin when PBMCs were stimulated overnight with Gag 263-272 (KK10, HLA-B*27⁺ peptide). Black horizontal bars indicate statistically significant difference (P < 0.05) between samples when stimulated overnight with the same variant peptide; gray horizontal bars indicate statistically significant difference (P < 0.05) between samples when stimulated overnight with opposite variant peptide.




Figure 5

Intracellular cytokine staining of CP CD8⁺ T cells. **A-D:** CD8⁺ T cells of HLA-B*57 positive CP were either freshly isolated (left) or stimulated with either wild type (center) or escape variant (right) HLA-B*57-restricted Gag peptides for 7 days. CPs were either on suppressive HAART regimens with undetectable viral loads (circles), were on HAART regimens but recently had detectable levels of viremia (triangles), or were not on HAART and had high levels of viremia (diamond). Cells from each group underwent an overnight stimulation with individual peptides. Percentage of CD8⁺ T cells expressing IFN-γ when restimulated with TW10 (**A**), KF11 (**B**), and IW9 (**C**) in blue, or the escape mutant variant peptide containing T242N/G248A (**A**), A163S (**B**), and I147L (**C**) mutations in red is shown. **E-H:** Percentage of CD8⁺ T cells expressing both IFN-γ and perforin after overnight stimulation with TW10 (**E**), KF11 (**F**), and IW9 (**G**) in blue, or the escape mutant variant peptide containing T242N/G248A (**E**), A163S (**F**), and I147L (**G**) mutations in red is shown. **D** and **H** show CD8⁺ T cells that express IFN-γ or co-express IFN-γ and perforin when PBMCs were stimulated overnight with Gag 263-272 (KK10, HLA-B*27⁺ peptide). Black asterisks indicate statistically significant difference (P < 0.05) between samples when restimulated with the same variant peptide; gray asterisks indicates statistically significant difference (P < 0.05) between samples when restimulated with opposite variant peptide.




Figure S1

Theoretical nonlinear regression curves plotted with each escape variant. Data generated in fitness assay is shown here (wild type, black; I147L, orange; A146P, navy; A163S, red; A146P/A163S, purple; T242N/G248A, brown; A146P/A163S/T242N/G248A, teal). Black line depicts theoretical nonlinear regression curve generated by GraphPad. R^2 values for each theoretical curve is show on the bottom right of individual plots. Plot on the bottom right depicts all theoretical curves corresponding to NL4-3 variant color on the same plot.




Figure S2

Comparison of stimulation status for suppressive function.

Suppressive capacity of CD8⁺ T cells, either unstimulated or stimulated with peptides corresponding to HLA-B*57 Gag epitope (WT) or escape mutant variant (Mutant), is compared for each of seven different escape mutant variant viruses used (wild type, black; I147L, orange; A146P, navy; A163S, red; A146P/A163S, purple; T242N/G248A, brown; A146P/A163S/T242N/G248A, teal).

Suppression on day 3 (left) and day 5 (right) is shown. n=7.

CHAPTER 2: ELITE SUPPRESSORS HAVE LOW LEVELS OF CELL-ASSOCIATED HIV-1 MRNA THAT INCREASE UPON T CELL ACTIVATION

Christopher W Pohlmeier*, C Korin Bullen*, Gregory M Laird*, Alyssa R Martin*, Stanley U Chioma, Victoria EK Walker-Sperling, and Joel N Blankson

*CPW, CKB, GML, and ARM contributed equally to this study

ABSTRACT

Objective:

Elite Controllers or Suppressors (ES) are patients who control HIV replication without antiretroviral therapy. In this study, we compared baseline and inducible HIV-1 mRNA levels in CD4+ T cells from ES and chronic progressors (CPs) receiving suppressive antiretroviral therapy.

Design/Methods:

We quantified basal levels of cell associated HIV-1 mRNA in CD4+ T cells isolated from CPs and ES. Additionally, we measured the fold upregulation of intracellular HIV-mRNA after stimulation of CD4+ T cells with phorbol 12-myristate 13-acetate (PMA) and ionomycin, and quantified the amount of HIV-mRNA levels released into culture supernatant.

Results:

ES have significantly less cell associated HIV-mRNA per 5×10^6 cells ($p = 0.003$); 8 of 10 CPs had quantifiable HIV-1 mRNA at baseline, whereas this was present in only 2 of 10 ES. Upon stimulation with PMA and ionomycin, 4 of 5 CPs and 7 of 9 ES showed increased cell associated HIV-mRNA. Interestingly, released HIV-1 mRNA could be detected in supernatants of CD4+ T cells stimulated with PMA/ionomycin from 5 of 8 ES.

Conclusion:

Our results demonstrate that while the baseline levels of cell associated HIV-1 mRNA are significantly lower in ES compared to CPs, stimulation of CD4+ T cells results in a comparable relative upregulation of viral transcription.

BACKGROUND

Elite controllers or suppressors (ES) are individuals infected with HIV that maintain undetectable viral loads without receiving antiretroviral therapy (ART). Prior studies have shown that replication-competent virus can be cultured from CD4⁺ T cells from some ES^{7,8,9,10,23,25,85} and full genome sequence analysis of replication competent virus has not revealed the presence of mutations associated with attenuation^{1,23}. Furthermore a recent study demonstrated that HIV-1 isolates from ES replicate vigorously and cause CD4⁺ T cell depletion in humanized mice²³. Low levels of HIV-1 RNA can be detected in the plasma of most ES with ultrasensitive PCR assays^{18,86,87,88}, and sequence analysis has revealed evolution of plasma virus present in these patients^{24,73,89,90,91}. These studies suggest that there is ongoing low-level replication in these patients, and yet in spite of this, ES maintain very small latent reservoirs as measured by total^{10,15,92} and integrated DNA⁹³ as well as by the quantitative viral outgrowth assay⁷ and the mouse viral outgrowth assay⁹⁴. Despite the small reservoirs present in these patients, we hypothesized that ES would have relatively high levels of cell associated mRNA as a result of ongoing viral replication.

Hatano and colleagues have measured cell-associated in PBMCs isolated from ES⁸⁸ and have compared cell associated RNA present in CD4⁺ T cells of ES and CPs isolated from gut-associated lymphoid tissue⁹⁵. These studies, however, did not specifically quantify HIV-1 mRNA which is a more direct indicator of HIV-1 transcription⁹⁶. In the present study, we measure baseline and inducible levels of cell associated HIV-mRNA in peripheral CD4⁺ T cells of ES and CPs using a previously validated primer probe set⁹⁷. These results deepen our understanding of the HIV-1 latent reservoir in ES.

RESULTS

Cell associated HIV-1 mRNA from unstimulated CD4+ T cells *ex vivo* was quantifiable in only 2 of 10 ES (median<10 copies/5M cells) as opposed to 8 of 10 CPs tested (median=92 copies/5M cells; $p=0.03$) (Figure 6a). Proviral HIV DNA was measured in 10 ES (median of 52 copies per 10^6 CD4+ T cells) and 10 CPs (median of 309 copies per 10^6 CD4+ T cells) to verify that our patients represent previously seen ES and CP cohorts (data not shown).

To determine whether transcription was induced with T cell activation, we compared cell associated HIV-1 mRNA at baseline and after maximal T cell activation. Treatment of CD4+ T cells with the T cell activation control of PMA plus ionomycin (PMA/I) increased cell associated HIV-1 mRNA levels in 7 of 9 ES and 4 of 5 CPs (Figure 6b). The measured cell associated HIV-1 mRNA per 10^6 CD4+ T cells remained statistically higher for CPs (median=33,676 per 10^6 cells) than for ES (median=873 per 10^6 cells) ($p=0.047$). However, there was no statistical difference between fold increase of cell associated HIV-mRNA between ES (13.1-fold, interquartile range=12.1) and CPs (19.2-fold, interquartile range=13.4, $p=0.22$) per 10^6 stimulated cells. Interestingly, cell associated HIV-1 mRNA levels remained below the limit of detection in 2 ES after stimulation with PMA and ionomycin stimulation.

We also assessed the effect of T cell activation on viral release into culture supernatant. Upon stimulation of replicates of 5×10^6 CD4+ T cells from ES with PMA and ionomycin, HIV-1 mRNA was detected in the culture supernatant from 5 of 8 ES (Figure 6c). 3 ES (ES6, ES9, and ES42) had undetectable HIV-1 mRNA measurements in culture supernatant even when a total of $10\text{--}20 \times 10^6$ CD4+ T cells were stimulated. In contrast, in ES22, ES23, and ES24 we detected HIV-1 mRNA in the culture supernatant from the majority of stimulated CD4+ T cell replicates. Interestingly in ES36 and ES43, HIV-1 mRNA was detected in the culture supernatant in only 1 of 5 stimulated replicates, with

3,012 and 5,116 copies of HIV-1 mRNA present in the positive wells. The frequency of detected released HIV-1 RNA suggests an 89% probability by Poisson distribution that the signal observed reflects a single cell releasing virus, and our HIV-1 mRNA measurements fit with our recent estimates of the burst size of an infected CD4+ T cell *in vitro*⁹⁸. Thus of the 25×10^6 cells stimulated, it appears that only 1 cell harbored an inducible virus in both of these patients. Interestingly, ES36 and ES43 also had undetectable cell associated HIV-1 mRNA in 5×10^6 CD4+ T cells after stimulation with PMA and ionomycin. Taken together, these data suggest that there is significant heterogeneity in the inducible HIV-1 reservoir seen in ES which is consistent with results obtained from the viral outgrowth assay^{7,23}.

DISCUSSION/CONCLUSION

Control of HIV-1 without ART by ES can be considered a model of a functional cure of infection. We have previously shown that these patients harbor much lower frequencies of latently infected CD4+ T cells⁷ than CPs on suppressive ART regimens⁹⁹, and in this study we further analyze the inducible reservoir in ES. We demonstrate that cell associated HIV-1 mRNA levels observed in ES CD4+ T cells at baseline were significantly lower than those observed in CPs. These results further characterize the nature the latent reservoir in ES patients recent studies that found that ES had low levels of HIV-1 RNA in peripheral blood and in the GALT^{88,95}. However, despite the low baseline levels of HIV-1 mRNA, there was no significant difference in the fold increase of cell associated HIV-1 mRNA levels upon PMA and ionomycin stimulation in CD4+ T cells isolated from ES and CPs, suggesting a similar induction of transcription in response to PMA and ionomycin treatment in both patient populations. While we cannot determine if this increase in HIV-1 mRNA is due to transcription of defective or intact proviral sequences¹⁰⁰, the data are consistent with our ability to culture replication-competent virus from some ES.

There was significant heterogeneity in the amount of cell associated and released virus seen in CD4+ T cells from different ES. Interestingly, ES23 and ES24 had detectable HIV-1 mRNA at baseline and both patients had a relatively high frequency of CD4+ T cells that released virus into culture supernatant following T cell stimulation. In contrast, the frequency of CD4+ T cells that released virus into culture supernatant was much lower in other ES. However we were able to culture and characterize replication-competent virus from 2 such patients, ES9⁹ and ES36²⁴ in prior studies. Further studies will be needed to determine whether virus that is released into culture supernatant following CD4+ T cell activation is in fact replication-competent.

Studies have shown that virus in the plasma of ES evolves over time suggesting that there is ongoing viral replication^{24,73,89,90,91}. In spite of that we still found very low levels of HIV-1 mRNA in peripheral CD4+ T cells in some of these patients. It is possible that the robust CTL response in these patients¹⁰¹ is able to eliminate transcriptionally active CD4+ T cells in peripheral blood that may be producing viral proteins, while CD4+ T cells in anatomical compartments that exclude CTL are the true source of ongoing viral replication. Results from a recent study that found that CD8+ T cells are excluded from B cell follicles in ES monkeys may be consistent with this hypothesis¹⁰².

In summary, we characterize baseline cell associated HIV-1 mRNA and show a robust upregulation following T cell stimulation in some ES. The results will have implications for patients who are subjected to curative procedures but still have residual low level HIV-1 reservoirs.

MATERIALS AND METHODS

Study Participants:

All studies were approved by the Johns Hopkins Institutional Review Board. All patients provided written informed consent before participation in this study. CPs are patients

with undetectable viral loads on suppressive ART regimens for at least 1 year. ES have maintained undetectable viral loads by standard commercial assays without ART.

Isolation of HIV-1 mRNA and DNA:

PBMCs were isolated from fresh blood samples by Ficoll gradient centrifugation. CD4⁺ T Cells were isolated using Miltenyi CD4⁺ T Cell Isolation Kit and cultured in RPMI1640 supplemented with 1% Pen/Strep and 10% FBS. 5x10⁶ cells in 2 mL of complete media were either cultured in the presence of DMSO (0.2%) or PMA (50 ng/mL) and Ionomycin (2 µM) for 24 hours. This stimulation resulted in greater than 90% of T cell activation as determined by CD69 expression. Supernatants were mixed with Trizol LS and cells were lysed using Trizol. Total RNA was isolated as described previously⁹⁶. Human genomic DNA was isolated using Puregene kit (Qiagen) as per manufacturer's instructions.

Quantification of HIV-1 mRNA and DNA:

Total intracellular RNA was reverse transcribed using qScript from Quanta Biosciences as per manufacturer's instructions. Reverse transcribed HIV-1 mRNA was quantified as described previously⁹⁶ using a primer and probe set that anneals to a highly conserved region of the 3' end of HIV-1 mRNA⁹⁷. The limit of quantification was set as the dilution point at which the Ct of the plasmid molecular standard replicates had an s.d. > 0.5. We determined that the limit of quantification for all intracellular HIV-mRNA transcripts was 10 copies⁹⁵. HIV-1 proviral DNA levels were quantified by qPCR using ToughMix (Quanta Biosciences). Previously published primers that target a small region of Gag¹⁰³ were used to detect HIV-1 proviral DNA, and cellular input was quantified measuring RNaseP levels (ThermoFisher) of a human genome standard (Roche).

FIGURES




Figure 6

HIV-mRNA before and after PMA and ionomycin stimulation. A

HIV-mRNA isolated from CD4⁺ T cells of ES (blue triangles) and CPs (red triangles) was quantified by qPCR. Samples below the limit of detection are represented by open triangles. The black horizontal bar represents the median value of HIV-mRNA for each population. HIV-mRNA levels observed were higher in CD4⁺ T cells of CPs than of ES ($p = 0.003$).

B CD4⁺ T cells of ES (blue inverted triangles) and CPs (red inverted triangles) were stimulated with either DMSO or PMA and ionomycin for 24 hours and HIV-mRNA was isolated and quantified by qPCR. Fold upregulation upon PMA and ionomycin stimulation is plotted. An empty symbol indicates the limit of detection was used as the DMSO control value. No significance difference in fold upregulation was detected between the two populations ($p = 0.27$ with empty symbol, 0.24 without empty symbol).

C HIV-mRNA release from CD4⁺ T cells upon PMA and ionomycin stimulation.

Replicates of 5×10^6 CD4⁺ T cells from ES were stimulated with PMA and ionomycin for 24 hours, and supernatant was collected for measurement of released HIV-mRNA. Open diamonds indicate replicates with supernatant values below the limit of detection. Single replicates from 2 ES (ES 36 and ES43) were positive, suggesting release of HIV-mRNA from individual cells.

SIGNIFICANCE AND FUTURE DIRECTIONS

The data presented here represent individual studies performed in a single lab. If I have learned anything in graduate school, I have learned to not believe any single study (or series of studies if done by the same lab) until the results are confirmed independently.

Nevertheless, the results discussed in this thesis suggest that some HIV-specific CD8+ T cells from ES are able to recognize both wild type and escape variant MHC-restricted epitopes. Without the ability to escape CTL control, viremia may be tempered. Activation and differentiation of naïve CD8+ T cells that express cross reactive T cell receptors may be critical targets for an effective therapeutic HIV vaccine.

While CTL activity is more robust in the ES population, this doesn't necessarily negate multiple factors being critical for the phenotype. Demonstrating that infected CD4+ T cells of ESs do not seem to have different levels of inducible intracellular HIV mRNA than CPs after maximal stimulation suggests that the resting state of latently infected CD4s is no different between the two populations. This supports the notion that more effective CTL control in CPs may lead to better control of viremia.

Understanding the *mechanism* by which ES control viremia may not be sufficient for developing a therapeutic vaccine. Cross-sectional studies cannot delineate between cause or consequence of the ES phenotype, i.e. whether any observed phenotype is causing the host to control viremia or is merely a byproduct of the host's control. It may be necessary to determine the *cause* of the ES phenotype in order to develop a therapeutic vaccine.

In that regard, it is best to study patients during seroconversion. This is very difficult to do, because a) ES are a very small proportion of the HIV-infected population and b) they will not necessarily present to a clinic during acute infection as most ES do not suffer

from acute retroviral syndrome. To circumvent this issue, Indian rhesus macaques are often used as ES models, though pathogenesis of SIV is often quite different than HIV; additionally, the rhesus macaque ES model may not truly recapitulate what is observed in humans.

Due to the enrichment of specific HLA molecules with the ES phenotype, studying seronegative donors that possess these protective alleles may shed some light on the cause of control. There is evidence to suggest that exposure to an antigen may induce an adaptive response against both the original antigen and HIV¹⁰⁴. Screening healthy donors for HIV-specific CD8⁺ T cells could demonstrate that the cause of immunological control is a pre-existing HIV-specific memory response. As finding an individual who would become an ES is rather challenging, inducing an HIV-specific phenotype in a humanized mouse model followed by HIV challenge and monitoring HIV viral load levels and CD8⁺ T cell activity may provide useful information on ways to induce the ES phenotype in individuals who would otherwise progress.

REFERENCES

- Okulicz JF, Lambotte O. Epidemiology and clinical characteristics of elite controllers. *Curr Opin HIV AIDS*. 2011, 6 (3): 163-168.
- O'Connell K a, Bailey JR, Blankson JN. Elucidating the elite: mechanisms of control in HIV-1 infection. *Trends Pharmacol Sci*. 2009, 30: 631-637.
- Deeks SG, Walker BD. Human immunodeficiency virus controllers: Mechanisms of durable virus control in the absence of antiretroviral therapy. *Immunity*. 2007, 27: 406-416.
- Migueles SA, Connors M. Long-term nonprogressive disease among untreated HIV-infected individuals: Clinical implications of understanding immune control of HIV. *JAMA*. 2010, 304: 194-201.
- Deacon NJ, Tsykin A, Solomon A, Smith K, Ludford-Mending M, Hooker DJ, McPhee DA, Greenway AL, Ellett A, Chatfield C, Lawson VA, Crowe S, Maerz A, Sonza S, Learmont J, Sullivan JS, Cunningham A, Dwyer D, Downton D, Mills J. Genomic structure of an attenuated quasi species of HIV-1 from a blood transfusion donor and recipients. *Science*. 1995, 270: 988-991.
- Alexander L, Weiskopf E, Greenough TC, Gaddis NC, Auerbach MR, Malim MH, O'Brien SJ, Walker BD, Sullivan JL, Desrosiers RC. Unusual polymorphisms in human immunodeficiency virus type 1 associated with nonprogressive infection. *J Virol*. 2000, 74: 4361-4376.
- Blankson JN, Bailey JR, Thayil S, Yang HC, Lassen K, Lai J, Gandhi SK, Siliciano JD, Williams TM, Siliciano RF. Isolation and characterization of replication-competent human immunodeficiency virus type 1 from a subset of elite suppressors. *J Virol*. 2007, 81 (5): 2508-2518.
- Lamine A, Caumont-Sarcos A, Chaix ML, Saez-Cirion A, Rouzioux C, Delfraissy JF, Pancino G, Lambotte O. Replication-competent HIV strains infect HIV controllers despite undetectable viremia (ANRS EP36 study). *AIDS*. 2007, 21 (8): 1043-1045.
- Bailey JR, O'Connell K, Yang HC, Han Y, Xu J, Jilek B, Williams TM, Ray SC, Siliciano RF, Blankson JN. Transmission of human immunodeficiency virus type 1 from a patient who developed AIDS to an elite suppressor. *J Virol*. 2008, 82 (15): 7395-7410.
- Julg B, Pereyra F, Buzon MJ, Piechocka-Trocha A, Clark MJ, Baker BM, Lian J, Miura T, Martinez-Picado J, Addo MM, Walker BD. Infrequent recovery of HIV from but robust exogenous infection of activated CD4(+) T cells in HIV elite controllers. *Clin Infect Dis*. 2010, 51 (2): 233-238. 10.1086/653677.
- Fraser C, Hollingsworth TD, Chapman R, de Wolf F, Hanage WP. Variation in HIV-1 set-point viral load: Epidemiological analysis and an evolutionary hypothesis. *PNAS*. 2007, 104 (44): 17441-17446.
- Kaslow R, Carrington M, Apple R, Park L. Influence of combinations of human major histocompatibility complex genes on the course of HIV-1 infection. *Nat Med*. 1996, 2 (4): 405-411.
- Vaidya SA, Streeck H, Beckwith N, Ghebremichael M, Pereyra F, Kwon DS, Addo MM, Rychert J, Routy JP, Jessen H, Kelleher AD, Hecht F, Sekaly RP, Carrington M, Walker BD, Allen TM, Rosenberg ES, Altfeld M. Temporal effect of HLA-B*57 on viral control during primary HIV-1 infection. *Retrovirology*. 2013, 10:139.
- Migueles SA, Sabbaghian MS, Shupert WL, Bettinotti MP, Marincola FM, Martino L, Hallahan CW, Selig SM, Schwartz D, Sullivan J, Connors M. HLA B*5701 is highly associated with restriction of virus replication in a subgroup of HIV-infected long term nonprogressors. *Proc Natl Acad Sci USA*. 2000, 97 (6): 2709-2714.
- Lambotte O, Boufassa F, Madec Y, Nguyen A, Goujard C, Meyer L, Rouzioux C, Venet A, Delfraissy JF, SEROCO-HEMOCO Study Group. HIV controllers: A homogeneous group of HIV-1-infected patients with spontaneous control of viral replication. *Clin Infect Dis*. 2005, 41 (7): 1053-1056.
- Emu B, Sinclair E, Hatano H, Ferre A, Shacklett B, Martin JN, McCune JM, Deeks SG. HLA class I-restricted T-cell responses may contribute to the control of human immunodeficiency virus infection, but such responses are not always necessary for long-term virus control. *J Virol*. 2008, 82 (11): 5398-5407.
- Han Y, Lai J, Barditch-Crovo P, Gallant JE, Williams TM, Siliciano RF, Blankson JN. The role of protective HCP5 and HLA-C associated polymorphisms in the control of HIV-1 replication in a subset of elite suppressors. *AIDS*. 2008, 22 (4): 541-544.
- Migueles SA, Osborne CM, Royce C, Compton AA, Joshi RP, Weeks KA, Rood JE, Berkley AM, Sacha JB, Coglianor-Shutta NA, Lloyd M, Roby G, Kwan R, McLaughlin M, Stallings S, Rehm C, O'Shea MA, Mican J, Packard BZ, Komoriya A, Palmer S, Wiegand AP, Maldarelli F, Coffin JM, Mellors JW, Hallahan CW, Follman DA, Connors M. Lytic granule loading of CD8+ T cells is required for HIV-infected cell elimination associated with immune control. *Immunity*. 2008, 29 (6): 1009-1021.
- Pereyra F, Addo MM, Kaufmann DE, Liu Y, Miura T, Rathod A, Baker B, Trocha A, Rosenberg R, Mackey E, Ueda P, Lu Z, Cohen D, Wrin T, Petropoulos CJ, Rosenberg ES, Walker BD. Genetic and immunologic heterogeneity among persons who control HIV infection in the absence of therapy. *J Infect Dis*. 2008, 197 (4): 563-571.
- Sajadi MM, Constantine NT, Mann DL, Charurat M, Dadzan E, Kadlecik P, Redfield RR. Epidemiologic characteristics and natural history of HIV-1 natural viral suppressors. *J Acquir Immune Defic Syndr*. 2009, 50 (4): 403-408.

21. Migueles SA, Laborico AC, Shupert WL, Sabbaghian MS, Rabin R, Hallahan CW, Van Baarle D, Kostense S, Miedema F, McLaughlin M, Ehler L, Metcalf J, Liu S, Connors M. HIV-specific CD8⁺ T cell proliferation is coupled to perforin expression and is maintained in nonprogressors. *Nat Immunol.* 2002, 3 (11): 1061-1068.
22. Betts MR, Nason MC, West SM, De Rosa SC, Migueles SA, Abraham J, Lederman MM, Benito JM, Goepfert PA, Connors M, Roederer M, Koup RA. HIV nonprogressors preferentially maintain highly functional HIV-specific CD8⁺ T cells. *Blood.* 2006, 107 (12): 4781-4789.
23. Salgado M, Swanson MD, Pohlmeier CW, Buckheit III RW, Wu J, Archin NM, *et al.* HLA-B*57 Elite Suppressor and Chronic Progressor HIV-1 Isolates Replicate Vigorously and Cause CD4⁺ T Cell Depletion in Humanized BLT Mice. *J Virol* 2014; 88:3340–52.
24. Bailey JR, Williams TM, Siliciano RF, Blankson JN. Maintenance of viral suppression in HIV-1-infected HLA-B*57⁺ elite suppressors despite CTL escape mutations. *J Exp Med.* 2006, 203 (5): 1357-1369
25. Thompson MA, Adberg JA, Hoy JF, Telenti A, Benson C, Cahn P, Eron JJ, Gunthard HF, Hammer SM, Reiss P, Richman DD, Rizzardini G, Thoma DL, Jacobsen DM, Volberding PA. Antiretroviral treatment of adult HIV infection: 2012 recommendations of the International Antiviral Society-USA panel. *JAMA.* 2012, 308(4):387-402.
26. Dinoso JB, Kim SY, Wiegand AM, Palmer SE, Gange SJ, Cranmer L, O'Shea A, Callender M, Spivak A, Brennan T, Kearney MF, Proschan MA, Mican JM, Rehm CA, Coffin JM, Mellors JW, Siliciano RF, Maldarelli F. Treatment intensification does not reduce residual HIV-1 viremia in patients on highly active antiretroviral therapy. *PNAS.* 2009, 106(23):9403-9408.
27. Casazza JP, Betts MR, Picker LJ, Koup RA. Decay kinetics of human immunodeficiency virus-specific CD8⁺ T cells in peripheral blood after initiation of highly active antiretroviral therapy. *J Virol.* 2001, 75(14):6508-6516.
28. Jolditch SJ, Eriksson EM, Tarosso LF, Kuebler PJ, Kallas EG, Nielsen EK, Wiznia AA, Rosenberg MG, Nixon DF. Decay kinetics of HIV-1 specific T cell responses in vertically HIV-1 exposed seronegative infants. *Front Immunol.* 2012, 2(94).
29. Buckheit RW, Allen TG, Alme A, Salgado M, O'Connell K a, Huculak S, Falade-Nwulia O, Williams TM, Gallant JE, Siliciano RF, Blankson JN: Host factors dictate control of viral replication in two HIV-1 controller/chronic progressor transmission pairs. *Nat Commun.* 2012, 3: 716
30. Itescu S, Mathur-Wagh U, Skovron M, Brancato L, Marmor M, Zeleniuch-Jacquotte A, Winchester R: HLA B35 is associated with accelerated progression to AIDS. *J Acquir Immune Defic Syndr.* 1991, 5 (1): 37-45.
31. Fellay J, Shianna KV, Ge D, Colombo S, Ledergerber B, Weale M, Zhang K, Gumbs C, Castagna A, Cossarizza A, Cozzi-Lepri A, De Luca A, Easterbrook P, Francioli P, Mallal S, Martinez-Picado J, Miro JM, Obel N, Smith JP, Wyniger J, Descombes P, Antonarakis SE, Letvin NL, McMichael AJ, Haynes BF, Telenti A, Goldstein DB: A whole-genome association study of major determinants for host control of HIV-1. *Science.* 2007, 317 (5840): 944-947. 10.1126/science.1143767.
32. Catano G, Kulkarni H, He W, Marconi VC, Agan BK, Landrum M, Anderson S, Delmar J, Telles V, Song L, Castiblanco J, Clark RA, Dolan MJ, Ahuja SK: HIV-1 disease-influencing effects associated with ZNRD1, HCP5 and HLA-C alleles are attributable mainly to either HLA-A10 or HLA-B*57 alleles. *PLoS One.* 2008, 3 (11): e3636-10.1371/journal.pone.0003636.
33. Dalmasso C, Carpentier W, Meyer L, Rouzioux C, Goujard C, Chaix ML, Lambotte O, Avettand-Fenoel V, Le Clerc S, de Senneville LD, Deveau C, Boufassa F, Debre P, Delfraissy JF, Broet P, Theodorou I, ANRS Genome Wide Association 01: Distinct genetic loci control plasma HIV-RNA and cellular HIV-DNA levels in HIV-1 infection: The ANRS genome wide association 01 study. *PLoS One.* 2008, 3 (12): e3907-10.1371/journal.pone.0003907.
34. Limou S, Le Clerc S, Coulonges C, Carpentier W, Dina C, Delaneau O, Labib T, Taing L, Sladek R, Deveau C, Ratsimandresy R, Montes M, Spadoni JL, Lelievre JD, Levy Y, Therwath A, Schachter F, Matsuda F, Gut I, Froguel P, Delfraissy JF, Hercberg S, Zagury JF, ANRS Genomic Group: Genomewide association study of an AIDS-nonprogression cohort emphasizes the role played by HLA genes (ANRS Genomewide Association Study 02). *J Infect Dis.* 2009, 199 (3): 419-426. 10.1086/596067.
35. van Manen D, Kootstra NA, Boeser-Nunnink B, Handulle MA, van't Wout AB, Schuitemaker H: Association of HLA-C and HCP5 gene regions with the clinical course of HIV-1 infection. *AIDS.* 2009, 23 (1): 19-28. 10.1097/QAD.0b013e32831db247.
36. International HIV Controllers S, Pereyra F, Jia X, McLaren PJ, Telenti A, de Bakker PI, Walker BD, Ripke S, Brumme CJ, Pulit SL, Carrington M, Kadie CM, Carlson JM, Heckerman D, Graham RR, Plenge RM, Deeks SG, Gianniny L, Crawford G, Sullivan J, Gonzalez E, Davies L, Camargo A, Moore JM, Beattie N, Gupta S, Crenshaw A, Burtt NP, Guiducci C, Gupta N, Gao X, *et al.* The major genetic determinants of HIV-1 control affect HLA class I peptide presentation. *Science.* 2010, 330 (6010): 1551-1557.

37. Almeida JR, Price DA, Papagno L, Arkoub ZA, Sauce D, Bornstein E, Asher TE, Samri A, Schnuriger A, Theodorou I, Costagliola D, Rouzioux C, Agut H, Marcelin AG, Douek D, Autran B, Appay V: Superior control of HIV-1 replication by CD8⁺ T cells is reflected by their avidity, polyfunctionality, and clonal turnover. *J Exp Med*. 2007, 204 (10): 2473-2485. 10.1084/jem.20070784.
38. Ferre AL, Hunt PW, Critchfield JW, Young DH, Morris MM, Garcia JC, Pollard RB, Yee HF, Martin JN, Deeks SG, Shackett BL: Mucosal immune responses to HIV-1 in elite controllers: a potential correlate of immune control. *Blood*. 2009, 113 (17): 3978-3989. 10.1182/blood-2008-10-182709.
39. Hersperger AR, Pereyra F, Nason M, Demers K, Sheth P, Shin LY, Kovacs CM, Rodriguez B, Sieg SF, Teixeira-Johnson L, Gudonis D, Goepfert PA, Lederman MM, Frank I, Makedonas G, Kaul R, Walker BD, Betts MR: Perforin expression directly *ex vivo* by HIV-specific CD8 T-cells is a correlate of HIV elite control. *PLoS Pathog*. 2010, 6 (5): e1000917-10.1371/journal.ppat.1000917.
40. Sáez-Cirión A, Lacabartz C, Lambotte O, Versmisse P, Urrutia A, Boufassa F, Barré-Sinoussi F, Delfraissy JF, Sinet M, Pancino G, Venet A: HIV controllers exhibit potent CD8 T cell capacity to suppress HIV infection *ex vivo* and peculiar cytotoxic T lymphocyte activation phenotype. *Proc Natl Acad Sci U S A*. 2007, 104 (16): 6776-6781. 10.1073/pnas.0611244104.
41. Crawford H, Lumm W, Leslie A, Schaefer M, Boeras D, Prado JG, Tang J, Farmer P, Ndugu'u T, Lakhi S, Gilmour J, Goepfert P, Walker BD, Kaslow R, Mulenga J, Allen S, Goulder PJR, Hunter E: Evolution of HLA-B*5703 HIV-1 escape mutations in HLA-B*5703-positive individuals and their transmission recipients. *J Exp Med*. 2009, 206: 909-921.
42. Bailey JR, Zhang H, Wegweiser BW, Yang H, Herrera L, Ahonkhai A, Williams TM, Siliciano RF, Blankson JN: Evolution of HIV-1 in an HLA-B * 57 – positive patient during virologic escape. *J Infect Dis*. 2007, 196: 50-55. 10.1086/518515.
43. Durand CM, Connell KAO, Apuzzo LG, Langan SJ, Ahonkhai AA, Ceccato CM, Williams TM, Margolick JB, Blankson JN: HIV-1 Gag evolution in recently infected HLA-B*57 patients with Low level viremia. *AIDS*. 2010, 24: 2405-2408.
44. Goonetilleke N, Liu MKP, Salazar-Gonzalez JF, Ferrari G, Giorgi E, Ganusov VV, Keele BF, Learn GH, Turnbull EL, Salazar MG, Weinhold KJ, Moore S, Letvin N, Haynes BF, Cohen MS, Hraber P, Bhattacharya T, Borrow P, Perelson AS, Hahn BH, Shaw GM, Korber BT, McMichael AJ: The first T cell response to transmitted/founder virus contributes to the control of acute viremia in HIV-1 infection. *J Exp Med*. 2009, 206: 1253-1272. 10.1084/jem.20090365.
45. Migueles SA, Laborico AC, Imamichi H, Shupert WL, Royce C, McLaughlin M, Ehler L, Metcalf J, Liu S, Hallahan CW, Connors M: The differential ability of HLA B * 5701 + long-term nonprogressors and progressors to restrict human immunodeficiency virus replication is Not caused by loss of recognition of autologous viral gag sequences. *J Virol*. 2003, 77 (12): 6889-6898. 10.1128/JVI.77.12.6889-6898.2003.
46. Bailey JR, Brennan TP, O'Connell KA, Siliciano RF, Blankson JN: Evidence of CD8⁺ T-cell-mediated selective pressure on human immunodeficiency virus type 1 nef in HLA-B*57⁺ elite suppressors. *J Virol*. 2009, 83: 88-97. 10.1128/JVI.01958-08.
47. Friedrich TC, Dodds EJ, Yant LJ, Vojnov L, Rudersdorf R, Cullen C, Evans DT, Desrosiers RC, Mothé BR, Sidney J, Sette A, Kunstman K, Wolinsky S, Piatak M, Lifson J, Hughes AL, Wilson N, O'Connor DH, Watkins DI: Reversion of CTL escape-variant immunodeficiency viruses *in vivo*. *Nat Med*. 2004, 10: 275-281. 10.1038/nm998.
48. Peyerl FW, Bazick HS, Newberg MH, Barouch DH, Sodroski J, Letvin NL: Fitness costs limit viral escape from cytotoxic T lymphocytes at a structurally constrained epitope. *J Virol*. 2004, 78: 13091-13910.
49. Matano T, Kobayashi M, Igarashi H, Takeda A, Nakamura H, Kano M, Sugimoto C, Mori K, Iida A, Hirata T, Hasegawa M, Yuasa T, Miyazawa M, Takahashi Y, Yasunami M, Kimura A, O'Connor DH, Watkins DI, Nagai Y: Cytotoxic T lymphocyte-based control of simian immunodeficiency virus replication in a preclinical AIDS vaccine trial. *J Exp Med*. 2004, 199: 1709-1718. 10.1084/jem.20040432.
50. Fernandez CS, Stratov I, De Rose R, Walsh K, Dale CJ, Smith MZ, Agy MB, Hu S-I, Krebs KC, Watkins DI, O'Connor DH, Davenport MP, Kent SJ: Rapid viral escape at an immunodominant simian-human immunodeficiency virus cytotoxic T-lymphocyte epitope exacts a dramatic fitness cost. *J Virol*. 2005, 79: 5721-5731. 10.1128/JVI.79.9.5721-5731.2005.
51. Martinez-Picado J, Prado JG, Fry EE, Pfafferott K, Leslie A, Chetty S, Thobakgale C, Honeyborne I, Crawford H, Matthews P, Pillay T, Rousseau C, Mullins JI, Brander C, Walker BD, Stuart DI, Kiepiela P,

- Goulder PJ: Fitness cost of escape mutations in p24 Gag in association with control of human immunodeficiency virus type 1. *J Virol.* 2006, 80: 3617-3623. 10.1128/JVI.80.7.3617-3623.2006.
52. Boutwell CL, Rowley CF, Essex M: Reduced viral replication capacity of human immunodeficiency virus type 1 subtype C caused by cytotoxic-T-lymphocyte escape mutations in HLA-B57 epitopes of capsid protein. *J Virol.* 2009, 83: 2460-2468. 10.1128/JVI.01970-08.
 53. Meiraz A, Garber OG, Harari S, Hassin D, Berke G: Switch from perforin-expressing to perforin-deficient CD8(+) T cells accounts for two distinct types of effector cytotoxic T lymphocytes *in vivo*. *Immunology.* 2009, 128 (1): 69-82. 10.1111/j.1365-2567.2009.03072.x.
 54. Zuñiga R, Lucchetti A, Galvan P, Sanchez S, Sanchez C, Hernandez A, Sanchez H, Frahm N, Linde CH, Hewitt HS, Hildebrand W, Altfeld M, Allen TM, Walker BD, Korber BT, Leitner T, Sanchez J, Brander C: Relative dominance of Gag p24-specific cytotoxic T lymphocytes is associated with human immunodeficiency virus control. *J Virol.* 2006, 80: 3122-3125. 10.1128/JVI.80.6.3122-3125.2006.
 55. Honeyborne I, Prendergast A, Pereyra F, Leslie A, Crawford H, Payne R, Reddy S, Bishop K, Moodley E, Nair K, van der Stok M, McCarthy N, Rousseau CM, Addo M, Mullins JI, Brander C, Kiepiela P, Walker BD, Goulder PJR: Control of human immunodeficiency virus type 1 is associated with HLA-B*13 and targeting of multiple gag-specific CD8⁺ T-cell epitopes. *J Virol.* 2007, 81: 3667-3672. 10.1128/JVI.02689-06.
 56. Kiepiela P, Ngumbela K, Thobakgale C, Ramduth D, Honeyborne I, Moodley E, Reddy S, de Pierres C, Mncube Z, Mkhwanazi N, Bishop K, van der Stok M, Nair K, Khan N, Crawford H, Payne R, Leslie A, Prado J, Prendergast A, Frater J, McCarthy N, Brander C, Learn GH, Nickle D, Rousseau C, Coovadia H, Mullins JI, Heckerman D, Walker BD, Goulder P: CD8⁺ T-cell responses to different HIV proteins have discordant associations with viral load. *Nat Med.* 2007, 13: 46-53. 10.1038/nm1520.
 57. Sacha JB, Chung C, Rakasz EG, Spencer SP, Jonas AK, Bean AT, Lee W, Burwitz BJ, Stephany JJ, Loffredo JT, Allison DB, Adnan S, Hoji A, Wilson NA, Friedrich TC, Lifson JD, Yang OO, Watkins DI: Gag-specific CD8⁺ T lymphocytes recognize infected cells before AIDS-virus integration and viral protein expression. *J Immunol.* 2007, 178: 2746-2754.
 58. Kloverpris HN, Payne RP, Sacha JB, Rasaiyaah JT, Chen F, Takiguchi M, Yang OO, Towers GJ, Goulder P, Prado JG: Early antigen presentation of protective HIV-1 KF11Gag and KK10Gag epitopes from incoming viral particles facilitates rapid recognition of infected cells by specific CD8⁺ T cells. *J Virol.* 2013, 87: 2628-2638. 10.1128/JVI.02131-12.
 59. Buseyne F, Le Gall S, Boccaccio C, Abastado JP, Lifson JD, Arthur LO, Riviere Y, Heard JM, Schwartz O: MHC-I-restricted presentation of HIV-1 virion antigens without viral replication. *Nat Med.* 2001, 7: 344-349. 10.1038/85493.
 60. Buckheit RW, Siliciano RF, Blankson JN: Primary CD8⁺ T cells from elite suppressors effectively eliminate non-productively HIV-1 infected resting and activated CD4⁺ T cells. *Retrovirology.* 2013, 10: 68-10.1186/1742-4690-10-68.
 61. Graf EH, Pace MJ, Peterson B, Lynch LJ, Chukwulebe SB, Mexas AM, Shaheen F, Martin JN, Deeks SG, Connors M, Migueles S, O'Doherty U: Gag-positive reservoir cells are susceptible to HIV-specific cytotoxic T lymphocyte mediated clearance. *PLoS One.* 2013, 8: e71879-10.1371/journal.pone.0071879.
 62. Mudd P a, Ericson AJ, Burwitz BJ, Wilson N, O'Connor DH, Hughes AL, Watkins DI: Escape from CD8(+) T cell responses in Mamu-B*00801(+) macaques differentiates progressors from elite controllers. *J Immunol.* 2012, 188: 3364-3370. 10.4049/jimmunol.1102470.
 63. Miura T, Brockman M, Brumme ZL, Brumme CJ, Pereyra F, Trocha A, Block BL, Schneidewind A, Allen TM, Heckerman D, Walker BD: HLA-associated alterations in replication capacity of chimeric NL4-3 viruses carrying gag-protease from elite controllers of human immunodeficiency virus type 1. *J Virol.* 2009, 83: 140-149. 10.1128/JVI.01471-08.
 64. Schneidewind A, Brockman M a, Yang R, Adam RI, Li B, Le Gall S, Rinaldo CR, Craggs SL, Allgaier RL, Power K a, Kuntzen T, Tung C-S, LaBute MX, Mueller SM, Harrer T, McMichael AJ, Goulder PJR, Aiken C, Brander C, Kelleher AD, Allen TM: Escape from the dominant HLA-B27-restricted cytotoxic T-lymphocyte response in Gag is associated with a dramatic reduction in human immunodeficiency virus type 1 replication. *J Virol.* 2007, 81: 12382-12389. 10.1128/JVI.01543-07.
 65. Crawford H, Prado JG, Leslie A, Hué S, Honeyborne I, Reddy S, van der Stok M, Mncube Z, Brander C, Rousseau C, Mullins JI, Kaslow R, Goepfert P, Allen S, Hunter E, Mulenga J, Kiepiela P, Walker BD, Goulder PJR: Compensatory mutation partially restores fitness and delays reversion of escape mutation

- within the immunodominant HLA-B*5703-restricted Gag epitope in chronic human immunodeficiency virus type 1 infection. *J Virol.* 2007, 81: 8346-8351. 10.1128/JVI.00465-07.
66. Klenerman P, Zinkernagel RM: Original antigenic sin impairs cytotoxic T lymphocyte responses to viruses bearing variant epitopes. *Nature.* 1998, 394: 482-485. 10.1038/28860.
 67. Francis TJ: On the doctrine of original antigenic Sin. *Proc Am Philos Soc.* 1960, 104: 572-578.
 68. Allen TM, Yu XG, Kalife ET, Reyrol LL, Lichterfeld M, John M, Cheng M, Allgaier RL, Mui S, Frahm N, Alter G, Brown NV, Johnston MN, Rosenberg ES, Mallal SA, Brander C, Walker BD, Altfeld M: De novo generation of escape variant-specific CD8⁺ T-cell responses following cytotoxic T-lymphocyte escape in chronic human immunodeficiency virus type 1 infection. *J Virol.* 2005, 79: 12952-12960. 10.1128/JVI.79.20.12952-12960.2005.
 69. Goulder PJR, Altfeld M, Rosenberg ES, Nguyen T, Tang Y, Eldridge RL, Addo MM, He S, Mukherjee JS, Phillips MN, Bunce M, Kalams SA, Sekaly RP, Walker BD, Brander C: Substantial differences in specificity of HIV-specific cytotoxic T cells in acute and chronic HIV infection. *J Exp Med.* 2001, 193: 181-194. 10.1084/jem.193.2.181.
 70. Lichterfeld M, Kavanagh DG, Williams KL, Moza B, Mui SK, Miura T, Sivamurthy R, Allgaier R, Pereyra F, Trocha A, Feeney M, Gandhi RT, Rosenberg ES, Altfeld M, Allen TM, Allen R, Walker BD, Sundberg EJ, Yu XG: A viral CTL escape mutation leading to immunoglobulin-like transcript 4-mediated functional inhibition of myelomonocytic cells. *J Exp Med.* 2007, 204: 2813-2824. 10.1084/jem.20061865.
 71. Feeney ME, Tang Y, Pfaffert K, Roosevelt KA, Draenert R, Trocha A, Yu XG, Verrill C, Allen T, Moore C, Mallal S, Burchett S, McIntosh K, Pelton SI, St John MA, Hazra R, Klenerman P, Altfeld M, Walker BD, Goulder PJR: HIV-1 viral escape in infancy followed by emergence of a variant-specific CTL response. *J Immunol.* 2005, 174: 7524-7530.
 72. Miura T, Brockman MA, Schneidewint A, Lobritz M, Pereyra F, Rathod A, Block BL, Brumme ZL, Brumme CJ, Baker B, Rothchild AC, Li B, Trocha A, Cutrell E, Frahm N, Brander C, Toth I, Arts EJ, Allen TM, Walker BD: HLA-B57/B*5801 human immunodeficiency virus type 1 elite controllers select for rare Gag variants associated with reduced viral replication capacity and strong cytotoxic T-lymphocyte recognition. *J Virol.* 2009, 83: 2743-2755. 10.1128/JVI.02265-08.
 73. O'Connell K a, Brennan TP, Bailey JR, Ray SC, Siliciano RF, Blankson JN: Control of HIV-1 in elite suppressors despite ongoing replication and evolution in plasma virus. *J Virol.* 2010, 84: 7018-7028. 10.1128/JVI.00548-10.
 74. Betts MR, Ambrozak DR, Douek DC, Bonhoeffer S, Brenchley JM, Casazza JP, Koup RA, Picker LJ: Analysis of total human immunodeficiency virus (HIV)-specific CD4⁺ and CD8⁺ T-cell responses: Relationship to viral load in untreated HIV infection. *J Virol.* 2001, 75: 11983-11991. 10.1128/JVI.75.24.11983-11991.2001.
 75. Addo MM, Yu XG, Rathod A, Cohen D, Eldridge RL, Strick D, Johnston MN, Corcoran C, Wurcel AG, Fitzpatrick CA, Feeney ME, Rodriguez WR, Basgoz N, Draenert R, Stone DR, Brander C, Goulder PJR, Rosenberg ES, Altfeld M, Walker BD: Comprehensive epitope analysis of human immunodeficiency virus type 1 (HIV-1)-specific T-cell responses directed against the entire expressed HIV-1 genome demonstrate broadly directed responses, but no correlation to viral load. *J Virol.* 2003, 77: 2081-2092. 10.1128/JVI.77.3.2081-2092.2003.
 76. Valentine LE, Piaskowski SM, Rakasz EG, Henry NL, Wilson N, Watkins DI: Recognition of escape variants in ELISPOT does not always predict CD8⁺ T-cell recognition of simian immunodeficiency virus-infected cells expressing the same variant sequences. *J Virol.* 2008, 82: 575-581. 10.1128/JVI.00275-07.
 77. Loffredo JT, Burwitz BJ, Rakasz EG, Spencer SP, Stephany JJ, Vela JPG, Martin SR, Reed J, Piaskowski SM, Furlott J, Weisgrau KL, Rodrigues DS, Soma T, Napoé G, Friedrich TC, Wilson N a, Kallas EG, Watkins DI: The antiviral efficacy of simian immunodeficiency virus-specific CD8⁺ T cells is unrelated to epitope specificity and is abrogated by viral escape. *J Virol.* 2007, 81: 2624-2634. 10.1128/JVI.01912-06
 78. Sáez-Cirión A, Lacabartz C, Lambotte O, Versmisse P, Urrutia A, Boufassa F, Barré-Sinoussi F, Delfraissy J-F, Sinet M, Pancino G, Venet A: HIV controllers exhibit potent CD8 T cell capacity to suppress HIV infection *ex vivo* and peculiar cytotoxic T lymphocyte activation phenotype. *Proc Natl Acad Sci USA.* 2007, 104: 6776-6781. 10.1073/pnas.0611244104.
 79. Sáez-Cirión A, Sinet M, Shin SY, Urrutia A, Versmisse P, Lacabartz C, Boufassa F, Avettand-Fènoël V, Rouzioux C, Delfraissy J-F, Barré-Sinoussi F, Lambotte O, Venet A, Pancino G: Heterogeneity in HIV

- suppression by CD8 T cells from HIV controllers: Association with Gag-specific CD8 T cell responses. *J Immunol.* 2009, 182: 7828-7837. 10.4049/jimmunol.0803928.
80. Buckheit RW, Salgado M, Siliciano RF, Blankson JN: Inhibitory potential of subpopulations of CD8⁺ T cells in HIV-1-infected elite suppressors. *J Virol.* 2012, 86: 13679-13688. 10.1128/JVI.02439-12.
 81. O'Connell K a, Hegarty RW, Siliciano RF, Blankson JN: Viral suppression of multiple escape mutants by de novo CD8(+) T cell responses in a human immunodeficiency virus-1 infected elite suppressor. *Retrovirology.* 2011, 8: 63-10.1186/1742-4690-8-63.
 82. Zhang H, Zho Y, Alcock C, Kiefer T, Monie D, Siliciano J, Li Q, Pham P, Cofrancesco J, Persaud D, Siliciano RF: Novel single-cell-level phenotypic assay for residual drug susceptibility and reduced replication capacity of drug-resistant human immunodeficiency virus type 1. *J Virol.* 2004, 78 (4): 1718-1729. 10.1128/JVI.78.4.1718-1729.2004.
 83. O'Doherty U, Swiggard WJ, Malim MH: Human immunodeficiency virus type 1 spinoculation enhances infection through virus budding. *J Virol.* 2000, 74 (21): 10074-10080. 10.1128/JVI.74.21.10074-10080.2000.
 84. Shan L, Rabi SA, Laird GM, Eisele EE, Zhang H, Margolick JB, Siliciano RF: A novel assay for quantification of HIV-1 RNA. *J Virol.* 2013, 87 (110): 6521-6525.
 85. Gaillard S, Dinoso JB, Marsh JA, DeZern AE, O'Connell KA, Spivak AM, *et al.* Sustained elite suppression of replication competent HIV-1 in a patient treated with rituximab based chemotherapy. *J Clin Virol* 2011; **51**:195–198.
 86. Dinoso JB, Kim SY, Siliciano RF, Blankson JN. A Comparison of Viral Loads between HIV-1–Infected Elite Suppressors and Individuals Who Receive Suppressive Highly Active Antiretroviral Therapy. *Clin Infect Dis* 2008; **47**:102–104.
 87. Pereyra F, Palmer S, Miura T, Block BL, Wiegand A, Rothchild AC, *et al.* Persistent Low-Level Viremia in HIV-1 Elite Controllers and Relationship to Immunologic Parameters. *J Infect Dis* 2009; **200**:984–990.
 88. Hatano H, Delwart EL, Norris PJ, Lee T-H, Dunn-Williams J, Hunt PW, *et al.* Evidence for persistent low-level viremia in individuals who control human immunodeficiency virus in the absence of antiretroviral therapy. *J Virol* 2009; **83**:329–35.
 89. Salgado M, Brennan TP, O'Connell KA, Bailey JR, Ray SC, Siliciano RF, *et al.* Evolution of the HIV-1 nef gene in HLA-B*57 Positive Elite Suppressors. *Retrovirology* 2010; **7**. doi:10.1186/1742-4690-7-94
 90. Mens H, Kearney M, Wiegand A, Shao W, Schønning K, Gerstoft J, *et al.* HIV-1 Continues To Replicate and Evolve in Patients with Natural Control of HIV Infection. *J Virol* 2010; **84**:12971–12981.
 91. Salgado M, Gandhi SK, Buckheit RW, Berkenblit G V, Blankson JN. Evolution of an Attenuated HIV-1 Isolate in an Elite Suppressor. *AIDS Res Hum Retroviruses* 2014; **30**:284–288.
 92. Gulck E Van, Bracke L, Heyndrickx L, Coppens S, Atkinson D, Pasternak A, *et al.* Immune and Viral Correlates of “Secondary Viral Control” after Treatment Interruption in Chronically HIV- 1 Infected Patients. *PLoS One* 2012; **7**:e37792.
 93. Metcalf Pate K a., Pohlmeier CW, Walker-Sperling VE, Foote JB, Najjarro KM, Cryer CG, *et al.* A murine viral outgrowth assay to detect residual HIV-1 in patients with undetectable viral loads. *J Infect Dis* 2015; :1–10.
 94. Hatano H, Somsouk M, Sinclair E, Harvill K, Gilman L, Cohen M, *et al.* Comparison of HIV DNA and RNA in Gut-Associated Lymphoid Tissue of HIV-Infected Controllers and Non-controllers. *AIDS* 2013; **27**:2255–2260.
 95. Bullen CK, Laird GM, Durand CM, Siliciano JD, Siliciano RF. New ex vivo approaches distinguish effective and ineffective single agents for reversing HIV-1 latency in vivo. *Nat Med* 2014; **20**:425–9.
 96. Shan L, Rabi SA, Laird GM, Eisele EE, Zhang H, Margolick JB, *et al.* A Novel PCR Assay for Quantification of HIV-1 RNA. *J Virol* 2013; **87**:6521–6525.
 97. O'Connell K a, Rabi SA, Siliciano RF, Blankson JN. CD4+ T cells from elite suppressors are more susceptible to HIV-1 but produce fewer virions than cells from chronic progressors. *Proc Natl Acad Sci U S A* 2011; **108**:E689–98.
 98. Siliciano JD, Siliciano RF. *Enhanced Culture Assay for Detection and Quantitation of Latently Infected, Resting CD4+ T-Cells Carrying Replication-Competent Virus in HIV-1-Infected Individuals.* *Methods in Molecular Biology* 2005; **304**:3-15.
 99. Ho Y, Pollack R, Yong P, Siliciano RF.: Defective HIV-1 Proviruses Can Be Transcribed Upon Activation [Abstract]. *CROI* 2015; **392**.
 100. Migueles SA, Connors M. Success and failure of the cellular immune response against HIV-1. *Nat Immunol* 2015; **16**:563–570.
 101. Fukazawa Y, Lum R, Okoye AA, Park H, Matsuda K, Bae JY, *et al.* B cell follicle sanctuary permits persistent productive simian immunodeficiency virus infection in elite controllers. *Nat Med* 2015; **21**:132–139.

102. Palmer S, Wiegand AP, Maldarelli F, Bazmi H, Mican JM, Polis M, *et al.* New Real-Time Reverse Transcriptase-Initiated PCR Assay with Single-Copy Sensitivity for Human Immunodeficiency Virus Type 1 RNA in Plasma. *J Clin Microbiol* 2003; **41**:4531–4536.
103. Su LF, Kidd BA, Han A, Kotzin JJ, Davis MM. Virus-specific CD4(+) memory-phenotype T cells are abundant in unexposed adults. *Immunity*. 2013, 38(2):373-383.

CURRICULUM VITAE

The Johns Hopkins University School of Medicine

Christopher Pohlmeier

April 2016

Educational History

2011-2016	Johns Hopkins University School of Medicine Baltimore, MD Ph.D., Biochemical Cellular and Molecular Biology
2007-2011	Johns Hopkins University School of Arts and Sciences Baltimore, MD B.S. Molecular and Cellular Biology

Research Experience

Metcalfe Pate KA, **Pohlmeier CW** *et al.* A murine outgrowth assay to detect residual HIV-1 in patients with undetectable viral loads. *J Infect Dis.* Epub ahead of print (2015).

Salgado M, Swanson MD, **Pohlmeier CW**, *et al.* HLA-B*57 elite suppressor and chronic progressor HIV-1 isolates replicate vigorously and cause CD4⁺ T cell depletion in humanized BLT mice. *J Virol.* 88, 3340-3352 (2014).

Pohlmeier CW, Buckheit RW III, Siliciano RF, Blankson JN. CD8⁺ T cells from HLA-B*57 elite suppressors effectively suppress replication of HIV-1 escape mutants. *Retrovirology.* 10:152 (2013).

Buss J, Coltharp C, Huang T, **Pohlmeier C**, Wang SC, Hatem C, Xiao J. In vivo organization of the FtsZ-ring by ZapA and ZapB revealed by quantitative super-resolution microscopy. *Mol Microbiol.* 89, 1099-1120 (2013).

Phua SC, **Pohlmeier CW**, Inoue T. Rapidly Relocating molecules between organelles to manipulate small GTPase activity. *ACS Chem Biol.* 12, 1950-1955 (2012).

Ueno Tasuku, Falkenburger BH, **Pohlmeier CW**, Inoue T. Triggering actin comets versus membrane ruffles: distinctive effects of phosphoinositides on actin reorganization. *Sci Signal.* 20, ra87 (2011).

Umeda N, Ueno T, **Pohlmeier CW**, *et al.* A photocleavable rapamycin conjugate for spatiotemporal control of small GTPase activity. *J Am Chem Soc.* 133, 12-14 (2011).

Posters and Presentations

Pohlmeier CW *et al.* Measurements of Viral Transcription in Elite Suppressors CD4⁺ T Cells. Themed discussion panel member. *CROI 2015.*

Pohlmeier CW *et al.* Measurements of Viral Transcription in Elite Suppressors CD4⁺ T Cells. Poster. *CROI 2015.*

Pohlmeier CW *et al.* CD8⁺ T Cells from HLA-B*57 Elite Suppressors Effectively Suppress Replication of HIV-1 Escape Mutants. Poster. *CROI 2014.*